

COUNTY OF MORRIS
DEPARTMENT OF PUBLIC WORKS
MOSQUITO CONTROL DIVISION

P.O. Box 900
Morristown, New Jersey 07963-0900

Board of County Commissioners
Commissioner Director
John Krickus

Commissioner Deputy Director
Christine Myers

Douglas R. Cabana
Stephen H. Shaw
Deborah Smith
Tayfun Selen
Thomas J. Mastrangelo



County Administrator
Deena Leary

*Director of Public Works &
County Engineer*
Christopher J. Vitz, P.E.
973-285-6750
Fax: 973-539-3141
cvitz@co.morris.nj.us

*Director, Mosquito
Control Division*
Kristian McMorland
973-285-6450
kmcmorland@co.morris.nj.us

“Anvil®”

This sheet answers some basic questions about a mosquito control product in use in Morris County. Morris County Division of Mosquito Control, along with several other resources (listed at the end of this sheet), can provide more detailed information.

What is Anvil® and how is it used?

Anvil® is an imitation mimic of a naturally occurring insecticide that is produced by certain chrysanthemum flowers. It is an adulticide that is recommended for mosquito control in New Jersey by the Agricultural Experiment Station of Rutgers University. It contains the active ingredients of “**sumithrin and piperonyl butoxide.**” The U.S. Environmental Protection Agency’s (EPA) current evaluation considers **sumithrin and piperonyl butoxide**- containing products to be slightly toxic with minimal potential risk to people when used properly as part of a complete mosquito control program.

Anvil® applications are made with Ultra Low Volume (ULV) equipment, which put out very small drops of the material at very low levels. While habitat management and measures to control immature mosquitoes in water are the preferred routine approaches, the spraying of adult mosquitoes is called for when biting populations reach critical levels or when a disease organism is present in adult mosquitoes. The application of **Anvil®** will only be conducted if deemed absolutely necessary. A very fine mist is sprayed into the air since flying mosquitoes must directly contact the pesticide in order for it to be effective. **Anvil®** is not residual and breaks down rapidly in sunlight, minimizing the buildup of this product in the environment.

How can I avoid exposure to Anvil®?

Risk to the general public from the use of **Anvil®** is minimal. Avoiding exposure is always the safest course of action, particularly for populations that may be at higher risk such as pregnant women, children, the elderly and those with chronic illness. Any possible exposure risk can be reduced by following some common sense actions:

- Pay attention to notices about spraying found through newspaper, websites, automated telephone messages or distributed by municipal, county or state agencies.
- Plan your activities to limit time spent outside during times of possible pesticide treatments.
- Move children’s toys out of application areas.
- Move animals and their food and water dishes out of application areas.
- Stay away from application equipment, whether in use or not.
- Whenever possible, remain indoors with windows closed and with window air conditioners on non-vent

(closed to the outside air) and window fans turned off during spraying.

- Avoid direct contact with surfaces that are still wet from pesticide spraying. Do not allow children to play in areas that have been sprayed until they have completely dried (approximately one hour).
- If you must remain outdoors, avoid eye and skin contact with the spray. If you get spray in your eyes or on your skin, immediately flush and rinse with water.

What are the symptoms of exposure to Anvil®?

Irritation or sensitization sometimes occurs after exposure, causing an asthmatic condition or skin rash. The chance of experiencing these symptoms of exposure with proper use is low. You should contact your physician, other medical providers or the New Jersey Poison Information and Education System (NJPIES) at **1-800-222-1222** if you experience these symptoms following a pesticide spraying. Bring this sheet with you if you visit a physician or other medical provider.

How long will Anvil® last in the environment?

The **Anvil®** spray stays in the air for a short time until it lands on surfaces. **Sumithrin** has a low persistence and lasts no longer than 25 days in water and soil. **Sumithrin** breaks down faster in sunlight.

Where can I get more information on Anvil®?

The following are resources for more information regarding **Anvil®** and mosquito control in your area (unless otherwise noted, available during normal business hours):

For overall pesticide-specific information – 9:30am to 7:30pm:

National Pesticide Information Center

800-858-7378

<http://npic.orst.edu>

For pesticide health information & possible exposures – 24 hours:

New Jersey Poison Information & Education System

800-222-1222

<http://www.njpies.org>

For New Jersey pesticide regulation & misuse complaints:

NJDEP Pesticide Control Program

609-984-6507

<http://www.state.nj.us/dep/enforcement/pcp/index.html>

For Federal pesticide regulations:

USEPA Region 2 Office of Pesticide Programs

877-251-4575

<http://www.epa.gov/ebtpages/pesticides.html>

For state-wide mosquito control information:

NJDEP Office of Mosquito Control Coordination

609-777-3666

<http://www.state.nj.us/dep/mosquito>

For local mosquito control information:

Morris County Division of Mosquito Control

973-285-6450

<http://www.morrismosquito.org>

For mosquito control recommendations:

Rutgers University, Department of Entomology

848-932-9774

<http://www-rci.rutgers.edu/~insects/indexentnj.htm>

EPA:

732-321-4391

<http://www.epa.gov/pesticides/factsheets/pyrethroids4mosquitos.htm>

Clarke Mosquito Control (Manufacturer):

800-323-5727

http://www.cmosquito.com/product_detail.cfm?productid=5&searchbox=anvil