

COUNTY OF MORRIS
DEPARTMENT OF PUBLIC WORKS
MOSQUITO CONTROL DIVISION

P.O. Box 900
Morristown, New Jersey 07963-0900

Board of County Commissioners
Commissioner Director
John Krickus

Commissioner Deputy Director
Christine Myers

Douglas R. Cabana
Stephen H. Shaw
Deborah Smith
Tayfun Selen
Thomas J. Mastrangelo



County Administrator
Deena Leary

*Director of Public Works &
County Engineer*
Christopher J. Vitz, P.E.
973-285-6750
Fax: 973-539-3141
cvitz@co.morris.nj.us

*Director, Mosquito
Control Division*
Kristian McMorland
973-285-6450
kmcmorland@co.morris.nj.us

“Duet Dual-Action® Adulcicide”®

This sheet answers some basic questions about a mosquito control product in use in Morris County. Morris County Division of Mosquito Control, along with several other resources (listed at the end of this sheet), can provide more detailed information.

What is *Duet Dual-Action®* adulcicide and how is it used?

Duet Dual-Action® contains two pesticides called *Prallethrin and Sumithrin*, and a synergistic compound called *piperonyl butoxide* which increases the effectiveness of the pesticides. Prallethrin and Sumithrin are members of a category of pesticides called *pyrethroids*, which in turn are synthetic versions of pesticides produced by plants called *pyrethrins*. Pyrethroid/piperonyl butoxide mixtures have been recommended for Ultra-Low-Volume (ULV) mosquito control in New Jersey by Rutgers, The State University of New Jersey. The U.S. Environmental Protection Agency’s (EPA) current evaluation considers pyrethroid-containing products to be slightly toxic with minimal potential risk to people when used properly as part of an integrated mosquito control program.

This pyrethroid-containing product is used for the control of adult mosquitoes. While habitat management and measures to control immature mosquitoes in water are preferred and most used, the spraying of adult mosquitoes is called for when biting populations reach critical levels or when a disease organism is present in adult mosquitoes. A very fine mist is sprayed into the air since flying mosquitoes must directly contact the pesticide in order for it to be effective. The combination of the two pesticides has been shown to produce what the manufacturer calls ‘benign agitation’. In other words mosquitoes are agitated from a resting state to a non-biting flying state where they are more vulnerable to pesticide exposure. This makes *Duet Dual-Action®* adulcicide more effective against hard-to-control species like *Aedes albopictus* which typically rest during the evening hours when adulciding usually takes place.

How can I reduce my exposure to *Duet Dual-Action®*?

Because of the very small amounts of active ingredients released per acre, the risk to the general public from the use of pyrethroid-containing products is minimal. Avoiding exposure is always the safest course of action. Any possible exposure risk can be reduced by following some common sense actions:

- Pay attention to notices about spraying found through newspapers, websites, automated telephone messages or distributed by municipal, county or state agencies.
- Plan your activities to limit time spent outside during times of possible pesticide treatments.

Offices located at the Hanover Garage on 120 E. Hanover Avenue, Cedar Knolls, New Jersey 07927

- Move your pets, their food, and water dishes inside during ULV application. Also bring clothing and children's toys inside.
- Stay away from application equipment, whether in use or not.
- Whenever possible, remain indoors with windows closed and with window air conditioners on non-vent (closed to the outside air) and window fans turned off during spraying.
- Avoid direct contact with surfaces that are still wet from pesticide spraying. Do not allow children to play in areas that have been sprayed until they have completely dried (approximately one hour).
- If you must remain outdoors, avoid eye and skin contact with the spray. If you get spray in your eyes or on your skin, immediately flush and rinse with water.

What are the symptoms of exposure to *Duet Dual-Action*[®] ?

Symptoms of over-exposure can include irritation to skin and eyes, respiratory and nasal irritation, irritability to sound or touch, abnormal facial sensation, sensation of prickling, tingling or creeping of skin, numbness, headache, dizziness, nausea, vomiting, diarrhea, excessive salivation, and fatigue. The chance of experiencing these symptoms of over-exposure with proper use is low. You should contact your physician, other medical providers, or the New Jersey Poison Information and Education System (NJPIES) at **1-800-222-1222** if you experience these symptoms following a pesticide spraying.

How long will *Duet Dual-Action*[®] last in the environment?

Pyrethroids have a soil half-life of 12 days. They have an extremely low pesticide movement rating because they bind tightly to the soil. Pyrethroids are unstable in light and air. They rapidly degrade in sunlight at the soil surface and in water. Piperonyl butoxide has a soil half-life of approximately 4 days.

Where can I get more information on this adulticide?

The following are resources for more information regarding *Duet Dual-Action*[®] and mosquito control in your area (unless otherwise noted, available during normal business hours):

For overall pesticide-specific information – 9:30am to 7:30pm:
National Pesticide Information Center **800-858-7378**
<http://npic.orst.edu>

For pesticide health information & possible exposures – 24 hours:
New Jersey Poison Information & Education System **800-222-1222**
<http://www.njpies.org>

For New Jersey pesticide regulation & misuse complaints:
NJDEP Pesticide Control Program **609-984-6507**
<http://www.state.nj.us/dep/enforcement/pcp/>

For Federal pesticide regulations:

USEPA Region 2 Office of Pesticide Programs
<http://www.epa.gov/ebtpages/pesticides.html>

877-251-4575

For state-wide mosquito control information:
NJDEP Office of Mosquito Control Coordination
<http://www.state.nj.us/dep/mosquito>

609-777-3666

For local mosquito control information:
Morris County Division of Mosquito Control
<http://morrismosquito.org>

973-285-6450

For mosquito control recommendations:
Rutgers University, Department of Entomology
<http://www-rci.rutgers.edu/~insects>

848-932-9774

Spraying for adult mosquitoes is a last resort. Most mosquito control work goes on “behind the scenes”, using water management, fish, and products to control immature mosquitoes in the water where they begin their life cycle. Controlling adult mosquitoes is more difficult because they are spread out and moving.